

Heat conduction in one-dimensional Yukawa chains

Bambi Hu^{1,2}, Haibin Li^{1,3}, and Bai-Song Xie^{1,4*}

¹*Department of Physics and Centre for Nonlinear Studies,
Hong Kong Baptist University, Hong Kong, China*

²*Department of Physics, University of Houston, Houston Tx 77204-5506*

³*Zhejiang Institute of Modern Physics, Zhejiang University, Hangzhou 310027, China*

⁴*Institute of Low Energy Nuclear Physics, Beijing Normal University, Beijing 100875, China*

(Dated: February 2, 2008)

Heat conduction in one-dimensional Yukawa chains is investigated. It is shown numerically that it has the abnormal heat conduction which is proportional to the system size. Effects of asymmetric external potential, the modified Frenkel-Kontorova one, on the heat conduction of system are also studied. It is found that the asymmetric property of external potential can induce asymmetric thermal conductivity that can be used to be a effective thermal rectifier. In certain of system parameters the heat flux are significantly different for two opposite direction. One can parametrically control the heat flux through this system by changing the potential strength and width.

PACS numbers: 44.10.+i, 05.45.-a, 52.27.Lw

Recent years many interesting works have been addressed to the problem of heat conduction [1, 2, 3, 4, 5, 6, 7, 8, 9]. It is well known that heat conductivity of one dimensional (1D) lattice is an important problem which is related to microscopic foundation of the Fourier law. It also provides an abundant example for the microscopic origin of the macroscopic irreversibility phenomenon. Detailed review of the problem is presented in the recent paper [1].

Except of extensive numerical studies focusing on the validity of Fourier law $J = -\kappa dT/dx$ for many models such as the Fermi-Pasta-Ulam (FPU) model [2, 3], the ding-a-ling model [4, 5], the Frenkel-Kontorova (FK) model [6, 7] and the Lorenz-gas model [8] and so on, some other different and important problems have also attracted people attention. For example most recently Casati *et al.* [9] investigated the possibility to control the energy transport inside a nonlinear 1D chain connecting with two thermostats at different temperatures. They emphasized that controlling heat conduction by nonlinearity opens a new possibility to design a thermal rectifier, i.e., a lattice that carries heat preferentially in one direction.

In this Letter we investigate heat conduction problem for a new type of model wherein there is a Yukawa-Coulomb interaction potential. It arises from dust-grain plasma which is extensively studied recently due to its new features with dust crystal or collective waves [10, 11, 12, 13, 14, 15]. Our study shows that the heat conductivity is abnormal in absence of external field as $\kappa \propto N$ as in harmonic chains. Under FK field, however, it is normal which is similar to the cases of former studies but with harmonic or/and FPU interactions [6]. Surprisingly for the dust chains with real wake potential [12], modified FK field, the heat conduction exhibits a mix-

ture property of normal and abnormal behaviors. It is found that the asymmetric property of external wake potential can induce asymmetric thermal conductivity that can be used to be a thermal rectifier. In a certain of system parameters, by changing the strength and width of wake potential, the heat fluxes for two opposite direction can be different remarkably.

In a general form the Hamiltonian of the considered system is written as

$$H = \sum_i h_i, \quad h_i = \frac{p_i^2}{2} + V(x_{i-1}, x_i) + U(x_i) \quad (1)$$

where $V(x_{i-1}, x_i)$ stands for the interaction potential of the nearest-neighbor particles and $U(x_i)$ is a external on-site potential. For dust particles system we have the interaction potential (refer to [14])

$$V(r_{ij}) = \frac{q^2}{r_{ij}} \exp(-\frac{r_{ij}}{\lambda_d}) - \alpha \frac{q^2}{r_{ij}} \quad (2)$$

where α is the parameter that ranges from 0 to 1, λ_d is the plasma Debye length and q is the dust grain charges, while r_{ij} is the distance between particles i -th and j -th. For simplicity we have assumed that the masses m_d and charges of all the grains q are the same and constant, while for convenience we shall choose parameter α that makes the neighbor dust particle is located at the bottom of V (see more clear in the following). Note that the plasma electrons and ions do not appear explicitly in the simulations but their effects, which result in an attractive force to dust, are taken into account in the effective interaction potential as the second term of Eq.(2). On the other hand in dusty plasma we know that there exist wake potential acting the dusts along downstream [12]

$$U(r_{ti}) = \frac{qq_t}{r_{ti}} \frac{2}{1 - M^{-2}} \cos(\frac{r_{ti}}{\lambda_d \sqrt{M^2 - 1}}) \quad (3)$$

where q_t is the upstream test dust particle charge, $M > 1$ is the Mach number of ion flow and r_{ti} is the distance of

*Author to whom correspondence should be addressed.




FIG. 1: Temperature profiles for different external potentials of field-free (a), FK (b) and FK-modified (c). The imposed temperature are $T_+ = 0.12$ and $T_- = 0.08$. The parameters are chosen as $\gamma_1 = 1$ and $\gamma_2 = 0.3$ (b), 2 (c). Averages are carried over a time interval 10^7 by dropping a transient 10^6 .

i -th particle from the test particle. The potential (3) is named as the FK-modified one in this paper because it is different from FK potential with varying amplitude.

Let us turn to the side of the motion of system. A general approach is employed here that are used by many authors like Lepri *et al.* and Hu *et al.*, namely, two Nose-Hoover thermostats are put on the first and last particle, keeping the temperature at T_+ and T_- , respectively. For simplicity and convenience, firstly we need to normalize the basic physical quantities: the mass by the particle mass m_d , the space distance by the dust inter-particle distance a_d and the time by the $\sqrt{ma_d^3/q^2}$ respectively. Then the equations of motion for particles are

$$\ddot{x}_1 = -\xi_+ \dot{x}_1 + f_1 - f_2, \quad \xi_+ = \frac{\dot{x}_1^2}{T_+} - 1$$




FIG. 2: Scaling of the heat flux JN with the number of particles N in the Yukawa interaction chains for different external potentials of field-free (a), FK (b) and FK-modified (c). System parameters and others are same as in Fig.1

$$\ddot{x}_i = f_i - f_{i+1}, \quad i = 2, \dots, N-1 \quad (4)$$

$$\ddot{x}_N = -\xi_- \dot{x}_N + f_N - f_{N-1}, \quad \xi_- = \frac{\dot{x}_N^2}{T_-} - 1$$

where $f_i = -V'(x_i - x_{i-1}) - U'(x_i)$ is the force. Usually the fixed boundary conditions $x_0 \equiv \text{const}$ and $x_{N+1} \equiv \text{const}$ are assumed. Now the interaction potential in normalized form can be written

$$V(x_{i-1}, x_i) = (\exp[-\chi(x_i - x_{i-1})] - \alpha) / (x_i - x_{i-1}) \quad (5)$$

where parameter $\chi = a_d/\lambda_d$ is the ratio of inter-particle distance to the plasma Debye length. Many studies show that when χ is large the nearest neighbor interaction is




FIG. 3: Dependence of J_L/J_R , the heat flux ratio of opposite direction for the Yukawa-Coulomb model, on parameter γ_2 , the strength of the FK-modified on-site external potential. The imposed temperature are $T_+ = 0.12$ and $T_- = 0.08$ for J_R and $T_+ = 0.08$ and $T_- = 0.12$ for J_L . $\gamma_1 = 1$ is given.

appropriate instead of particle-pair interaction [15]. In our following numerical studies $\chi = 5$ is generally used. The particles are arranged initially at equilibrium positions of V and the normalized inter-distance equals to 1, therefore, it means that parameter α , adjusting the attraction of plasma to dust particle, must satisfy $\alpha = (1 + \chi) \exp(-\chi)$. For the external FK-modified potential Eq.(3) we have its normalized form

$$U(x_i) = \gamma_2 \cos(\gamma_1 x_i)/x_i, \quad (6)$$

where $\gamma_1 = \chi/\sqrt{M^2 - 1}$ and $\gamma_2 = 2\beta/(1 - M^{-2})$ with $\beta = q_t/q$ the ratio of upstream test particle charge to downstream particles charge. Now we denote x_i as the position of i -th dust particle in the chain from the origin of test dust particle $x_{0t} \equiv 0$. In correspondence for the normalized external FK potential one has $U(x_i) = \gamma_2 \cos(\gamma_1 x_i)$. The different temperature profiles for different external potentials of field-free, FK and FK-modified are shown in Fig.1. The imposed temperature are $T_+ = 0.12$ and $T_- = 0.08$ and $\gamma_1 = 1$. The potential strengths are $\gamma_2 = 0.3$ and 2 for FK and FK-modified, respectively. Obviously the temperature gradients are constructed globally only in the FK case.

Usually the heat flux at the i -th position is given by $J_i(t) = \frac{1}{2}(\dot{x}_i + \dot{x}_{i+1})f_{i+1}$. Numerically the time average $J = \langle J_i(t) \rangle$ is independent of the index for long time enough. We plot in Fig.2 the heat flux JN vs the system size N for three different cases of field-free (a), FK (b) and FK-modified (c). Obviously the abnormal heat conductivity $\kappa \propto N$ is obtained for purely Yukawa interaction chains, which is similar to the case of harmonic situation. In the Yukawa chains for standard FK external potential, although it seems unclear about whether there exists phase transition of heat conduction between normal and abnormal [6, 7], our results indicates that the normal heat conductivity exists, which support the conclusion of normal heat conduction in Ref. [6]. From Fig.2 (c), however, we can see that it is abnormal globally on heat conduction in the Yukawa chains under FK-modified on-site external potential while the local temperature gradient appeared in the left region of system, see Fig.1(c). The dependence of J_L/J_R , the heat flux ratio of opposite direction for the Yukawa-Coulomb model,




FIG. 4: Momentum excitations for different types of external potentials of field-free [(a), (b)], FK [(c), (d)] and FK-modified [(e), (f)]. $p_L(0)$ and $p_R(0)$ denote the initial momentum kick on the left 1-th particle and right 100-th particle, respectively. For (c)-(f) the parameters of $\gamma_1 = 1$ and $\gamma_2 = 2$ are given.

on parameter γ_2 , the strength of the FK-modified on-site external potential is plotted in Fig.3, where the power law $J_L/J_R \propto 10^{0.24\gamma_2}$ is fitted in our numerical parameter regime. Our numerical experiments shows the validity of this power law until to $\gamma_2 = 4$ which indicates that the difference of two directional heat current can possibly over one order. However in practice of dusty plasmas it can only be realized by a relatively small γ_2 due to the test dust charge equals or less than the charge of particles in chains. For example, for given $\chi = 5$, $\gamma_1 = 1$ and $\beta = q_t/q \approx 1$ it corresponds to the practice ion flow Mach number $M = \sqrt{26}$ (usually in the sheath $M > 1$), and $\gamma_2 \approx 2$ which indicates $J_L/J_R \approx 3$.

In order to see how the single kicked particle transfer energy to the others, in Fig.4 we plotted the momentum excitations with different external potentials of three cases of field-free, FK and FK-modified. We can see that in the case of field-free there exists a solitary-like wave which can carry energy independently and not transform it to others. In case of FK there exist phonons which can transfer heat to the others. And in the case of FK-modified there seems a combination of two kind waves mentioned above: solitary-like and scatter phonons. Moreover it seems that there are always more solitary-like waves which propagates in the left-direction than that in the right direction. For example, there are




FIG. 5: Dependence of the heat flux ratio of two cases, for the Yukawa chains with different inverse width of the FK-modified on-site external potential, on temperature T . $\gamma_2 = 2$ is given.

three solitons in Fig.3(f) but only one in Fig.3(e). Maybe this is one reason why in this case the left-directional heat flux, J_L , is larger than that of right directional, J_R .

On the other hand according to the expression of heat flux, the ratio of them of two-direction is approximately

$$\frac{J_L}{J_R} \approx \sqrt{\frac{T_+}{T_-}} \frac{r_>}{r_<} \exp \chi(r_> - r_<) \quad (7)$$

where $r_>$ and $r_<$ are the average inter-particle distance in the region where the temperature is almost constant for $[T_+, T_-]$, right-direction heat-flow J_R , and $[T_-, T_+]$, left-direction heat-flow, respectively. In general it is shown that $r_>$ is a little larger than $r_<$ by numerical experiments. It is not surprising that the more phonons are scattered in the case of $[T_+, T_-]$ than that of vice versa. Therefore even if $r_>$ is just a very little greater than $r_<$, J_L/J_R would be greatly larger than one. For example, $r_> \approx 1.08$ and $r_< \approx 0.92$, then we have $J_L/J_R \approx 3$. However in the case of Harmonic interaction potential because of its symmetric quadratic property about equilibrium position the ratio of opposite directional heat flux is in order of $\sqrt{T_+/T_-}$, for example by choosing $T_+ = 0.12$,

$T_- = 0.08$, the ratio of $J_L/J_R \approx 1.23$. Our numerical results confirm this simple approximation analysis. Indeed if we instead of harmonic interaction chains the heat flux difference for two direction can not be over 23% under asymmetric potential.

Finally we calculate heat flux in different potential width γ_1^{-1} (see Fig.5). It shows that we can also control the energy flow by changing the potential width $\gamma_1^{-1} = \sqrt{M^2 - 1}/\chi$ through adjusting the ion flow velocity, which is associated with M or and the plasma particles density which is associated with χ . Obviously the great difference of heat flux at low temperature caused by the width of potential smear out as the temperature increases to over 0.5. It is noted that the ratio of heat flux $J_R(\gamma_1 = 1)/J_R(\gamma_1 = \pi)$ is about over two orders difference.

In summary we have studied the heat conduction problem in the system of 1D Yukawa interaction chains with asymmetric on-site modified FK wake potential. We can parametrically control the heat flux through this system by changing the potential strength and width. Moreover it is also possible to adjust the heat flux by changing the potential from FK to modified FK in the Yukawa chains, thus we provide a simple method to change the properties of the system, from a normal conductor obeying Fourier law, down to an almost perfect insulator. Some theoretical consideration has also been attempted to understand for our results and it is found that in the case of Yukawa interaction potential, the asymmetric property about equilibrium position has played a crucial role to improve the ratio of opposite directional heat flux. It seems that there exist some modes which are combinations of solitary-like and scattering phonons. Certainly more formidable theoretical works is still challenging our research on this problem in the future.

Authors would like to thank Drs. L. Wang, L. Yang and H. Zhang for their useful discussions. This work was supported by the grants of the Hong Kong Research Grants Council (RGC) and the Hong Kong Baptist University Faculty Research Grant (FRG). B. S. X also acknowledge the support of funding in part by the National Natural Science Foundation of China (NSFC) under Grant Nos. 10275007 and 10135010.

[1] S. Lepri, R. Livi, and A. Politi, Phys. Rep. **377**, 1-80 (2003).
[2] S. Lepri, R. Livi, and A. Politi, Phys. Rev. Lett. **78**, 1896 (1997).
[3] B. Hu, B. Li, and H. Zhao, Phys. Rev. E **61**, 3828 (2000).
[4] G. Casati, J. Ford, F. Vivaldi, and W. M. Visscher, Phys. Rev. Lett. **52**, 1861 (1984).
[5] H. A. Posch and Wm. G. Hoover, Phys. Rev. E **58**, 4344 (1998).
[6] B. Hu, B. Li, and H. Zhao, Phys. Rev. E **57**, 2992 (1998).
[7] A. V. Savin, and O. V. Gendelman, Phys. Rev. E **67**,

041205 (2003).
[8] D. Alonso, R. Artuso, G. Casati, and I. Guarneri, Phys. Rev. Lett. **82**, 1859 (1999).
[9] M. Terraneo, M. Peyrard and G. Casati, Phys. Rev. Lett. **88**, 094302 (2002).
[10] J. H. Chu and Lin I, Phys. Rev. Lett. **72**, 4009 (1994).
[11] H. Thomas et al., Phys. Rev. Lett. **73**, 652 (1994).
[12] S. V. Vladimirov and M. Nambu, Phys. Rev. E **52**, R2172 (1995).
[13] B. S. Xie and M. Y. Yu, Phys. Rev. E **62**, 8501 (2000).
[14] B. S. Xie and Z. A. Yang, Phys. Plasmas **9**, 4851 (2002).

- [15] D. Samsonov, A. V. Ivlev, R. A. Quinn and G. Morfill,
Phys. Rev. Lett. **88**, 095004 (2002).